レジメン名　：　GS（GEM＋TS-1）療法

疾患名　：　胆道がん

投与間隔（1コース期間）　：21日間

総コース数　：　PDになるまで

特記事項　：

	投与順序	レジメン内容（医薬品名）	投与量	単位（mg/㎡,V等）	投与方法（手技）	投与時間（速度等）	投与日(Day1等)	備考（内服薬の用法等）
Rp.	1	生理食塩液　100mL	1	本	点滴静注		D1、D8	ルート確保用
		生理食塩液　50mL	1	本				
	2	ﾃﾞｷｻｰﾄ　6.6mg	1	瓶	点滴静注	15分	D1、D8	
		ゲムシタビン	1000	mg/㎡				
	3	5%ブドウ糖液　100mL	1	本	点滴静注	30分	D1、D8	
	4	生理食塩液　50mL	1	本	点滴静注	15分	D1、D8	
		TS-1　60～120mg/日　14日間内服　(2投1休)						

減量基準

減量規定（副作用名等）	減量条件（検査値等）	抗癌剤名	減量割合等（減量後の割合又は投与量）

参考文献

・Kanai M, et al. A multi-institution phase II study of gemcitabine/S-1 combination chemotherapy for patients with advanced biliary tract cancer. Cancer Chemother Pharmacol. 2011; 67; 1429-1434

・Sasaki T, et al. Multicenter, phase II study of gemcitabine and S-1 combination chemotherapy in patients with advanced biliary tract cancer. Cancer Chemother Pharmacol. 2010; 65; 1101-1107

③Morizane C, et al. Randomized phase II study of gemcitabine plus S-1 versus S-1 in advanced biliary tract cancer: a Japan Clinical Oncology Group trial (JCOG 0805). Cancer Sci, 2013; 104; 1211-1216

・Sasaki T, et al. A randomized phase II study of gemcitabine and S-1 combination therapy versus gemcitabine monotherapy for advanced biliary tract cancer. Cancer Chemother Pharmacol. 2013; 71; 973-979