レジメン名　：　CPT-11+ベクティビックス

疾患名　：　大腸がん（KRAS野生型）

投与間隔（1コース期間）　：　14日間

総コース数　：　PDになるまで

特記事項　：　フィルター付きルートの使用
UGT1A1の測定、KRAS測定

	投与順序	レジメン内容（医薬品名）	投与量	単位（mg/㎡, V等）	投与方法（手技）	投与時間（速度等）	投与日（Day1等）	備考（内服薬の用法等）
Rp.	1	生理食塩液　100mL	1	本	点滴静注		D1	ルート確保用/フィルター付きルートの使用
		ﾊﾟﾛﾉｾﾄﾛﾝ点滴静注0.75mg	1	袋				
		ﾃﾞｷｻｰﾄ　6.6mg	1	瓶				
	2	ﾃﾞｷｻｰﾄ　1.65mg	2	本	点滴静注	15分	D1	
		ベクティビックス	6	mg/kg				
	3	生理食塩液　100mL	1	本	点滴静注	60分	D1	
	4	生理食塩液　50mL	1	本	点滴静注	15分	D1	
		イリノテカン	150	mg/㎡				
	5	生理食塩液　250mL	1	袋	点滴静注	90分	D1	
	6	生理食塩液　50mL	1	本	点滴静注	15分	D1	

減量基準

減量規定（副作用名等）	減量条件（検査値等）	抗癌剤名	減量割合等（減量後の割合又は投与量）

参考文献

・Seymour MT, et al. Panitumumab and irinotecan versus irinotecan alone for patients with KRAS wild-type, fluorouracil-resistant advanced colorectal cancer (PICCOLO): a prospectively stratified randomised trial. Lancet oncol. 2013; 14; 749-59

・Peeters M, et al. Randomized phase III study of panitumumab with fluorouracil, leucovorin, and irinotecan (FOLFIRI) compared with FOLFIRI alone as second-line treatment in patients with metastatic colorectal cancer. J Clin Oncol. 2010; 28; 4706-4713

・Ander T, et al. Panitumumab combined with irinotecan for patients with KRAS wild-type metastatic colorectal cancer refractory to standard chemotherapy: a GERCOR efficacy, tolerance, and translational molecular study. Ann oncol. 2013; 24; 412-9