レジメン名　：　XELOX

疾患名　：　大腸がん

投与間隔（1コース期間）　：21日毎

総コース数　：　術後補助　3ヶ月または6ヶ月
進行・再発　PDになるまで

特記事項　：

	投与順序	レジメン内容（医薬品名）	投与量	単位（mg/㎡,V等）	投与方法（手技）	投与時間（速度等）	投与日(Day1等)	備考（内服薬の用法等）
Rp.	1	生理食塩液　100mL	1	本	点滴静注		D1	ルート確保用
		ﾊﾟﾛﾉｾﾄﾛﾝ点滴静注0.75m	1	袋				
	2	デキサート　6.6mg	1	瓶	点滴静注	15分	D1	
		オキサリプラチン	130	mg/㎡				
	3	5%ブドウ糖液　500mL	1	袋	点滴静注	120分	D1	
	4	生理食塩液　50mL	1	本	点滴静注	15分	D1	
		ゼローダ　2000mg/㎡/日　2投1休						

減量基準

減量規定（副作用名等）	減量条件（検査値等）	抗癌剤名	減量割合等（減量後の割合又は投与量）
有害事象	Grade 3以上	オキサリプラチン	100mg/㎡　→85mg/㎡

参考文献

・Cassidy j, rt al. Randomized phase III study of capecitabine plus oxaliplatin compared with fluorouracil/folinic acid plus oxaliplatin as first-line therapy for metastatic colorectal cancer. J Clin Oncol. 2008; 26; 2006-12

②Rothenberg ML, et al. Capecitabine plus oxaliplatin (XELOX) versus 5-fluorouracil/folinic acid plus oxaliplatin (FOLFOX-4) as second-line therapy in metastatic colorectal cancer: a randomized phase III noninferiority study. Ann of Oncol, 2008; 19; 1720-6

③Doi T, et al. Phase I/II study of capecitabine plus oxaliplatin (XELOX) plus bevacizumab as first-line therapy in Japanese patients with metastatic colorectal cancer. Jpn J Clin Oncol. 2010; 40; 913-20