レジメン名　：　AC療法

疾患名　：　乳がん

投与間隔（1コース期間）　：　21日間

総コース数　：　術前術後療法4コース

特記事項　：　ケモセーフ使用

	投与順序	レジメン内容（医薬品名）	投与量	単位（mg/㎡,V等）	投与方法（手技）	投与時間（速度等）	投与日(Day1等)	備考（内服薬の用法等）
Rp.	1	生理食塩液　100mL	1	本	点滴静注		D1	ルート確保用/ケモセーフ使用/ｱﾌﾟﾚﾋﾟﾀﾝﾄ125mg内服確認すること
	2	ﾊﾟﾛﾉｾﾄﾛﾝ点滴静注0.75m	1	袋				
		デキサート　6.6mg	1	瓶				
		デキサート　1.65mg	2	本	点滴静注	15分	D1	
	3	ドキソルビシン	60	mg/㎡				生食50mL抜き取る
		生理食塩液　100mL	50	mL	点滴静注	30分	D1	
	4	エンドキサン	600	mg/㎡				
		生理食塩液　100mL	1	本	点滴静注	30分	D1	
	5	生理食塩液　50mL	1	本	点滴静注	15分	D1	

減量基準

減量規定（副作用名等）	減量条件（検査値等）	抗癌剤名	減量割合等（減量後の割合又は投与量）

参考文献

・Fisher B, rt al. Two months of doxorubicin-cyclophosphamide with and without interval reinduction therapy compared with 6 months of cyclophosphamide, methotrexate, and fluorouracil in positive-node breast cancer patients with tamoxifen-nonresponsive tumors: results from the National Surgical Adjuvant Breast and Bowel Project B-15. J Clin Oncol 1990; 8: 1483-96.

・Fisher B, rt al.Tamoxifen and chemotherapy for axillary node-negative, estrogen receptor-negative breast cancer: findings from National Surgical Adjuvant Breast and Bowel Project B-23. J Clin Oncol 2001; 19(4) : 931-42.